herpy

SAPOS-FLECHA

## Blue-bellied Poison Frog

# *Andinobates minutus*

*Andinobates minutus* is one of the smallest poison dart frogs, only 13-16 mm from snout to vent. It lives in the leaf litter of humid lowland forest from central Panama to the Pacific coast of Colombia, and the male carries the tadpoles up into bromeliads. The species is not collected for the pet trade and is hardly kept, so little is known about its care.

Família: Dendrobatidae  Nome anterior: *Ranitomeya minuta; Dendrobates minutus; Minyobates minutus*

### DADOS PRINCIPAIS

#### O ANIMAL

| | |
|---|---|
| Tamanho adulto | **13-16 mm**<br>comprimento focinho-cloaca |
| Longevidade | **5-10 anos** |
| Ativo | **De dia** |
| Estilo de vida | **Terrestre** |
| Habitat | **Floresta tropical** |
| Toxinas na pele na natureza | **Leves** |
| Toxinas na pele em cativeiro | **Nenhuma**<br>Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo |
| Dificuldade | **Experiente** |

#### CUIDADOS

| | |
|---|---|
| Terrário mín. | **45 × 45 × 45 cm**<br>C × L × A para 2 adultos |
| Temperatura, dia | **22-25 °C** |
| Temperatura, noite | **19-22 °C** |
| Umidade | **80-100%** |
| Água | **Sem recipiente de água** |
| UVB | **Opcional**<br>Zona de Ferguson 1 |
| Alimentação | **Pequenos invertebrados (colêmbolos, drosófilas)** |
| Mantido | **Em casal** |

#### REPRODUÇÃO

| | |
|---|---|
| Reprodução | **Ovíparo** |
| Desenvolvimento | **Girinos carregados até a água** |
| Local de postura | **No chão** |
| Ovos por postura | **1-8** |
| Fase larval | **60-80 dias**<br>da eclosão até a metamorfose |
| Alimentação das larvas | **Ração completa** |
| Cuidado parental | **Carrega os girinos** |

#### PROTEÇÃO E DOCUMENTAÇÃO

| | |
|---|---|
| CITES | **Apêndice II** |
| IUCN | **Menos preocupante (LC)** |

## Resumo dos cuidados

| | |
|---|---|
| **O TERRÁRIO** | Little is known about the care of *Andinobates minutus*. The care is based on the habitat: warm, humid lowland forest with rain for most of the year and a thick layer of leaf litter. |
| **TEMPERATURA** | The daytime temperature should be 22-25 °C and the night 19-22 °C. The temperature must not exceed 27 °C for long, and above 29 °C is dangerous, so the enclosure must not stand in sun or close to a heat source. |
| **UMIDADE E NEBULIZAÇÃO** | Humidity should be 80-100 %. Spray the enclosure 1-2 times a day with rainwater or reverse osmosis water so that the leaf litter is kept damp, or use a rain system on a timer. |
| **ÁGUA E QUALIDADE DA ÁGUA** | The adult frogs do not need a water bowl; they take up water through the skin from spraying and the damp leaf litter. |
| **ILUMINAÇÃO E UVB** | The species is diurnal and lives in shade on the forest floor. LED lighting for about 12 hours a day is enough for the plants. UVB is not necessary; a weak UVB light in Ferguson zone 1 can be used if there is plenty of shade. |
| **ALIMENTAÇÃO** | In the wild *Andinobates minutus* lives on mites, ants, springtails and insect larvae. |
| **COMPORTAMENTO E MANUSEIO** | The species is diurnal and stays in the leaf litter and by the hiding places. The males call from fixed spots and defend their territory, so a pair is the way to keep them. |
| **ÁREA DE OCORRÊNCIA NATURAL** | Colômbia e Panamá. |