herpy

SAPOS-FLECHA

Andean Poison Frog

Andinobates opisthomelas

Andinobates opisthomelas is a small, diurnal poison dart frog of 15-20 mm from the Andes of Colombia, where it lives in the leaf litter of humid mountain forest in Antioquia and Caldas. The species is Vulnerable in the wild, and almost nothing is known about its care.

Família: Dendrobatidae Nome anterior: *Ranitomeya opisthomelas; Dendrobates opisthomelas; Minyobates opisthomelas*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**15-20 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**1-2**
Eclosão	**16-24 dias**
Fase larval	**108-112 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Carrega os girinos**

CUIDADOS

Terrário mín.	**45 × 50 × 60 cm** C × L × A para 5 adultos
Temperatura, dia	**22-25 °C**
Temperatura, noite	**18-21 °C**
Umidade	**80-100%**
Água	**Sem recipiente de água**
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	Almost nothing is known about the care of *Andinobates opisthomelas*, and the species is hardly kept. The care is based on the habitat, the cool and humid mountain forest with leaf litter and bromeliads, and on close relatives.
TEMPERATURA	The daytime temperature should be 22-25 °C, and at night it may fall to 18-21 °C as in the mountain forest. Keep the temperature below 27 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 % and never below 70 %. Spray twice a day, morning and afternoon, or use a fogger or rain system with reverse osmosis water or rainwater.
ÁGUA E QUALIDADE DA ÁGUA	A water bowl is not needed; the frogs drink droplets from plants and spray water. For the tadpoles, use bromeliads with water in the leaf axils or small plastic cups with 2-3 cm of water.
ILUMINAÇÃO E UVB	The light is primarily for the plants: plant lighting for 10-12 hours a day provides a daily rhythm and growth.
ALIMENTAÇÃO	The food is small arthropods: springtails, small fruit flies (*Drosophila melanogaster*) and small isopods. Newly metamorphosed froglets take only springtails.
COMPORTAMENTO E MANUSEIO	*Andinobates opisthomelas* is diurnal and lives on the ground, so it stays low in the enclosure, in the leaf litter and by the bromeliads.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia.