herpy

SAPOS-FLECHA

Santander Poison Frog

Andinobates virolinensis

Andinobates virolinensis is a small, diurnal poison dart frog of up to 19 mm with a scarlet head and forebody and a brown hindbody. It is found only in the cloud forest of the eastern Andes in Colombia. The species is vulnerable in the wild and is hardly ever kept, so its care is based on the cool, humid mountain climate of its habitat.

Família: Dendrobatidae Nome anterior: *Ranitomeya virolinensis; Dendrobates virolinensis; Minyobates virolinensis*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**19 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**45 × 45 × 60 cm** C × L × A para 3 adultos
Temperatura, dia	**19-23 °C**
Temperatura, noite	**15-19 °C**
Umidade	**85-100%**
Água	**Sem recipiente de água**
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**1-2**
Eclosão	**16-24 dias**
Fase larval	**108-152 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Andinobates virolinensis*, and the species is hardly ever kept in captivity. Its care is based on the habitat: cool, humid cloud forest with a thick layer of leaf litter and bromeliads.
TEMPERATURA	The temperature has not been measured in the species' habitat. 19-23 °C by day and 15-19 °C at night suit cloud forest at 1,300-2,400 m.
UMIDADE E NEBULIZAÇÃO	Humidity should be 85-100 % with damp leaf litter all year round.
ÁGUA E QUALIDADE DA ÁGUA	The species does not need a water section. The water is found in the leaf axils of the bromeliads, where the tadpoles grow up, and it must be changed regularly so that it does not turn cloudy.
ILUMINAÇÃO E UVB	*A. virolinensis* lives on the shaded forest floor, so the light is primarily for the plants. Provide 10-12 hours of light a day with LED or fluorescent tubes and darkness at night. It is not documented that the species needs UVB.
ALIMENTAÇÃO	The diet is small arthropods: springtails and fruit flies (*Drosophila*) in the smallest size, and small woodlice. Feed 3-4 times a week, and always keep live feeder insects in the enclosure.
COMPORTAMENTO E MANUSEIO	Behaviour in captivity is not documented. In the wild several adults share the same bromeliad, so the species presumably tolerates company and can be kept as a pair or a small group.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia.