herpy

SAPOS-FLECHA

Anthony's Poison Arrow Frog

Epipedobates anthonyi

Epipedobates anthonyi is a small, red and striped poison frog of 19-27 mm from south-western Ecuador and north-western Peru. It is the frog that has been sold for decades as "tricolor", and it is probably the most widely kept and bred poison frog of all: hardy, bold, vocal and a prolific breeder.

Família: Dendrobatidae Nome anterior: *Dendrobates anthonyi; Phyllobates anthonyi*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**19-27 mm** comprimento focinho-cloaca
Longevidade	**5-14 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta seca**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**5-10 meses**
Ovos por postura	**10-40**
Eclosão	**9-14 dias** a 22-24 °C
Fase larval	**43-89 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 3 adultos
Temperatura, dia	**22-26 °C**
Temperatura, noite	**18-21 °C**
Umidade	**80-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Quase ameaçado (NT)**

Resumo dos cuidados

O TERRÁRIO	A terrarium of at least 40 × 40 × 40 cm suits one male and two females of *Epipedobates anthonyi*. A group of 5-10 animals needs at least 60 × 60 × 40 cm, and more animals need more floor area.
TEMPERATURA	Keep 22-26 °C by day with both cooler and warmer zones, and let the temperature fall to room temperature at night, around 18-21 °C. The temperature must not exceed 27 °C for long, and above 29 °C is dangerous.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 %. The species tolerates down to 50 % for short periods if it has access to water, but low humidity without water can quickly become fatal.
ÁGUA E QUALIDADE DA ÁGUA	Make one or more small water holes with about 2 cm of water, e.g. saucers, drinking dishes or half Brazil nut shells that the frogs can easily climb out of.
ILUMINAÇÃO E UVB	The light is for the plants. Fluorescent tubes or LED bars for about 12 hours a day are enough, and the day length may vary a little over the year.
ALIMENTAÇÃO	*Epipedobates anthonyi* is small but greedy and takes relatively large prey.
COMPORTAMENTO E MANUSEIO	*Epipedobates anthonyi* is diurnal, bold and often out in the open, and the males call with a trill, mostly in the morning and after misting.
ÁREA DE OCORRÊNCIA NATURAL	Equador e Peru.