herpy

SAPOS-FLECHA

Marbled Poison Frog

Epipedobates boulengeri

Epipedobates boulengeri is a small, diurnal poison dart frog of about 17-24 mm with a dark brown back and an unbroken, pale stripe along the side. It lives in the lowland rainforest along the Pacific coast from south-western Colombia to north-westernmost Ecuador. The species is hardly ever kept, and little is known about its care.

Família: Dendrobatidae Nome anterior: *Dendrobates boulengeri; Phyllobates boulengeri*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**17-24 mm** comprimento focinho-cloaca
Longevidade	**8-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**10-30**
Eclosão	**9-14 dias**
Fase larval	**42-70 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 3 adultos
Temperatura, dia	**22-26 °C**
Temperatura, noite	**19-22 °C**
Umidade	**80-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Epipedobates boulengeri*, and the species is hardly ever kept.
TEMPERATURA	Keep 22-26 °C during the day with slightly cooler and slightly warmer spots in the enclosure, and 19-22 °C at night. This is normally achieved with room temperature and the lighting; heat mats are unnecessary.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 %. Spray several times a day or use a rain system, for example three times a day for 20-60 seconds, preferably supplemented with a fogger.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water bowl with 1-2 cm of water, for example a low saucer or half a Brazil nut shell, gives the frogs a place to drink and bathe and is a breeding site when the male is to deposit tadpoles.
ILUMINAÇÃO E UVB	The light is for the plants: fluorescent tubes or LED bars for about 12 hours a day. The closely related species thrive without UVB, so it is optional.
ALIMENTAÇÃO	The frogs are given live, small prey: springtails and fruit flies (*Drosophila melanogaster*), and adults also take the larger *Drosophila hydei* and newly hatched crickets.
COMPORTAMENTO E MANUSEIO	The species of the genus are diurnal, ground-dwelling and territorial, and the males call from fixed spots. The call has not been described in more detail for *E. boulengeri* in the narrow sense.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia e Equador.