herpy

SAPOS-FLECHA

Darwin Wallace poison frog

Epipedobates darwinwallacei

Epipedobates darwinwallacei is a small, black poison dart frog of 13-20 mm with orange to yellow spots. It lives in the montane forest around Mindo in north-western Ecuador and is threatened. Since 2024 it has been regarded as a synonym of *Epipedobates espinosai*. Little is known about its care.

Família: Dendrobatidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**13-20 mm** comprimento focinho-cloaca
Longevidade	**8-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical e floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 3 adultos
Temperatura, dia	**20-24 °C**
Temperatura, noite	**18-20 °C**
Umidade	**80-100%**
Água	**Área aquática** profundidade 2-5 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**10-30**
Eclosão	**9-14 dias**
Fase larval	**42-70 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo (EN)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Epipedobates darwinwallacei*, and the species is hardly ever kept.
TEMPERATURA	Keep 20-24 °C during the day and 18-20 °C at night. This corresponds to the cool end of what suits *Epipedobates tricolor* from the same elevation. It is normally achieved with room temperature and the lighting.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 %. Spray several times a day or use a rain system, for example three times a day for 20-60 seconds, preferably supplemented with a fogger.
ÁGUA E QUALIDADE DA ÁGUA	Provide a small, permanent water feature with 2-5 cm of water and a gentle slope or ramp so that the frogs can easily climb out.
ILUMINAÇÃO E UVB	The light is for the plants: fluorescent tubes or LED bars for about 12 hours a day. The closely related species thrive without UVB, so it is optional.
ALIMENTAÇÃO	The frogs are given live, small prey: springtails and fruit flies (*Drosophila melanogaster*), and adults also take the larger *Drosophila hydei* and newly hatched crickets.
COMPORTAMENTO E MANUSEIO	*E. darwinwallacei* is diurnal and moves openly between grass and leaf litter. The call is very similar to the call of the brown *E. espinosai* from the lowlands. The males of the genus are territorial and call from fixed spots.
ÁREA DE OCORRÊNCIA NATURAL	Equador.