herpy

SAPOS-FLECHA

Phantasmal poison frog

Epipedobates tricolor

Epipedobates tricolor is a small, dark brown poison frog of 19-23 mm with yellow to yellow-green stripes and spots. It occurs only on the western slopes of the Andes in central Ecuador and is threatened. Almost all "tricolor" in the trade are in fact the closely related *Epipedobates anthonyi*; true *E. tricolor* is rare in terrariums.

Família: Dendrobatidae Nome anterior: *Dendrobates tricolor; Phyllobates tricolor*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**19-23 mm** comprimento focinho-cloaca
Longevidade	**5-14 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical e floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**5-10 meses**
Ovos por postura	**10-20**
Eclosão	**9-14 dias** a 22-24 °C
Fase larval	**42-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 3 adultos
Temperatura, dia	**22-26 °C**
Temperatura, noite	**18-20 °C**
Umidade	**80-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	One male and two females of *Epipedobates tricolor* need an enclosure of at least 40 × 40 × 40 cm. A group of 5-10 animals needs at least 60 × 60 × 40 cm, and more animals require more floor area.
TEMPERATURA	Keep 22-26 °C during the day with both cooler and warmer zones, and let the temperature drop to 18-20 °C at night.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 %. The species tolerates down to 50 % for short periods if it has access to water, but low humidity without water can quickly become fatal.
ÁGUA E QUALIDADE DA ÁGUA	Make one or more small water holes with about 2 cm of water, e.g. saucers, drinking dishes or half Brazil nut shells that the frogs can easily climb out of. The male may use them to deposit tadpoles.
ILUMINAÇÃO E UVB	The lighting is for the plants. Fluorescent tubes or LED bars for about 12 hours a day are enough, and the day length can vary slightly over the year.
ALIMENTAÇÃO	*Epipedobates tricolor* is small but greedy and takes relatively large prey.
COMPORTAMENTO E MANUSEIO	The males of *Epipedobates tricolor* call with a soft, trilling voice, mostly in the morning and especially after misting, most often from an elevated perch.
ÁREA DE OCORRÊNCIA NATURAL	Equador.