herpy

SAPOS-FLECHA

Rio Santiago Poison Frog

Excidobates captivus

Excidobates captivus is a very small, diurnal poison frog of 12-17 mm from the border region between Peru and Ecuador. It is black with orange-red spots in rows along the back and yellow spots on the limbs and belly, and it lives on the forest floor in humid rainforest and montane forest.

Família: Dendrobatidae Nome anterior: *Dendrobates captivus; Ranitomeya captiva; Adelphobates captivus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**12-17 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical e floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**22-26 °C**
Temperatura, noite	**18-22 °C**
Umidade	**80-100%**
Água	**Sem recipiente de água**
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**4-8**
Fase larval	**65-75 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Excidobates captivus*. The species lives on the forest floor in warm to mild, permanently humid forest.
TEMPERATURA	The temperature at the species' localities has not been measured. 22-26 °C by day and 18-22 °C at night lies between the climate in the lowlands and at 800-1200 m elevation.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 %, and the substrate should be damp all year. Mist with reverse-osmosis or rainwater once or twice a day, or use a rain or fogging system. Make sure the air does not stand still.
ÁGUA E QUALIDADE DA ÁGUA	A water dish is not necessary, because the frogs get moisture from the misting and the damp leaf litter. The water in bromeliads and film canisters is replaced with fresh reverse-osmosis or rainwater every week.
ILUMINAÇÃO E UVB	The species lives in shade on the forest floor, so the lighting should first and foremost support plant growth and a day of 12 hours. UVB is not necessary, but optional.
ALIMENTAÇÃO	*Excidobates captivus* is so small that it can only take very small feeder insects: springtails, flightless fruit flies (*Drosophila melanogaster*) and the smallest woodlice.
COMPORTAMENTO E MANUSEIO	The species is diurnal, but hides a great deal in the foliage. The male calls short, shrill sounds from hidden spots. It is best kept as a pair, because several animals together can cause conflict.
ÁREA DE OCORRÊNCIA NATURAL	Peru e Equador.