herpy

SAPOS-FLECHA

Trinidad Poison Frog

Mannophryne trinitatis

Mannophryne trinitatis is a small, brown, diurnal frog of 25-28 mm that occurs only on Trinidad, where it lives by clear, stony streams in humid forest. The male can turn jet black when calling, and the female has a clear yellow throat. The species is hardly ever kept, and little is known about its care.

Família: Aromobatidae Nome anterior: *Phyllobates trinitatis*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**25-28 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical e área úmida**
Toxinas na pele	**Nenhuma conhecida**
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**60 × 40 × 45 cm** C × L × A para 2 adultos
Temperatura, dia	**23-27 °C**
Temperatura, noite	**20-22 °C**
Umidade	**80-100%**
Água	**Área aquática** profundidade 3-5 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**6-13**
Eclosão	**21 dias**
Fase larval	**40-50 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about keeping *Mannophryne trinitatis*.
TEMPERATURA	The daytime temperature should be 23-27 °C and the night 20-22 °C. Frogs from forest streams tolerate heat poorly, so keep the temperature below 28 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 %. Mist daily so that leaves and stones are wet. A fogging system makes it easier to keep the humidity stable. Provide ventilation so that the air does not become stagnant.
ÁGUA E QUALIDADE DA ÁGUA	The species needs a permanent water section with shallow, clean and preferably gently flowing water, 3-5 cm deep, with stones and leaves.
ILUMINAÇÃO E UVB	UVB is hardly necessary, but a low dose (Ferguson zone 1) is optional. Plant lighting for 10-12 hours keeps moss and ferns alive. Provide shade with plants and bark so that the frogs can move away from the bright light.
ALIMENTAÇÃO	The species needs small arthropods: small and large fruit flies (*Drosophila melanogaster* and *Drosophila hydei*), springtails and small isopods.
COMPORTAMENTO E MANUSEIO	*Mannophryne trinitatis* is diurnal and territorial. Females defend their spots more often than males, with threats, wrestling and chasing, and they display the yellow throat as a threat.
ÁREA DE OCORRÊNCIA NATURAL	Trinidad e Tobago.