herpy

SAPOS-FLECHA

Polkadot Poison Frog

Oophaga arborea

Oophaga arborea is a small poison dart frog of 20-22 mm that lives high up in the trees in the humid montane forest of western Panama. It is black or brown with round yellow spots, breeds in bromeliads 10-16 m above the ground, and the tadpoles feed only on eggs from the female. The species is critically endangered.

Família: Dendrobatidae Nome anterior: *Dendrobates arboreus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**20-22 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola**
Habitat	**Floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Especialista**

CUIDADOS

Terrário mín.	**60 × 45 × 80 cm** C × L × A para 2 adultos
Temperatura, dia	**20-24 °C**
Temperatura, noite	**15-19 °C**
Umidade	**70-100%**
Água	**Sem recipiente de água**
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Nas axilas das folhas (por exemplo, bromélias)**
Ovos por postura	**4-8**
Eclosão	**14 dias**
Fase larval	**40-50 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ovos não fertilizados da fêmea**
Cuidado parental	**Vigia os ovos, carrega os girinos e alimenta os girinos com ovos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo crítico (CR)**

Resumo dos cuidados

O TERRÁRIO	Almost nothing is known about the care of *Oophaga arborea*, and the species is almost unknown in captivity.
TEMPERATURA	The daytime temperature should be 20-24 °C, and at night it should fall by at least 4-5 °C, to about 15-19 °C. This corresponds to the climate of the cloud forest with an annual mean temperature of about 19 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-80 % by day and rise to 100 % at night. Use plenty of mist and rain, and let the enclosure dry quickly afterwards with the help of a fan; stagnant, wet air is dangerous.
ÁGUA E QUALIDADE DA ÁGUA	A water bowl is not necessary. The frogs get moisture from mist, rain and the water-filled bromeliads, which are also used for the tadpoles.
ILUMINAÇÃO E UVB	The species lives high up in the canopy, where the light is stronger than on the forest floor, but the mist dampens it.
ALIMENTAÇÃO	Adults are fed small, flightless fruit flies, springtails, newly hatched micro crickets and field plankton; the same feeder insects are used for the other species of the genus.
COMPORTAMENTO E MANUSEIO	*Oophaga arborea* is a diurnal tree frog that is rarely seen on the ground.
ÁREA DE OCORRÊNCIA NATURAL	Panamá.