herpy

SAPOS-FLECHA

Granular Poison Frog

Oophaga granulifera

Oophaga granulifera is a small, diurnal poison dart frog of 19-22 mm from the Pacific side of Costa Rica and the far west of Panama. The skin is distinctly granular, the back red, orange or olive-green, and the legs and belly blue-green to turquoise. The species lives along shaded forest streams, and the female feeds her tadpoles with unfertilised eggs.

Família: Dendrobatidae Nome anterior: *Dendrobates granuliferus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**19-22 mm** comprimento focinho-cloaca
Longevidade	**6 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**80 × 40 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**21-25 °C**
Temperatura, noite	**17-21 °C**
Umidade	**80-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**12-18 meses**
Ovos por postura	**2-5**
Eclosão	**19-20 dias**
Fase larval	**87-212 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ovos não fertilizados da fêmea**
Cuidado parental	**Vigia os ovos, carrega os girinos e alimenta os girinos com ovos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the husbandry of *Oophaga granulifera*, and the species is rare in the hobby.
TEMPERATURA	The daytime temperature should be 21-25 °C, and the frogs are most active at 20-22 °C. At night the temperature should fall by 4-5 °C, to about 17-21 °C. Seasonal variation is minimal, about 1-2 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-90 %, and 100 % early in the morning and late in the day when the enclosure is misted.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish of 1-2 cm depth, for example a low bowl sunk into the substrate or a water section between the stones, provides drinking water and humidity.
ILUMINAÇÃO E UVB	The species lives on the shady forest floor, so the light must first and foremost cover the plants' needs.
ALIMENTAÇÃO	Adults are fed small, flightless fruit flies, springtails, newly hatched micro crickets, the smallest waxworm larvae and field plankton.
COMPORTAMENTO E MANUSEIO	*Oophaga granulifera* is shy and lives very secretively, especially in the dry season, when it almost completely shuts down.
ÁREA DE OCORRÊNCIA NATURAL	Costa Rica e Panamá.