herpy

SAPOS-FLECHA

Harlequin Poison Frog

Oophaga histrionica

Oophaga histrionica is a diurnal poison frog of 3-4 cm from the rainforest of western Colombia, brown to black with large orange spots. The tadpoles live exclusively on infertile eggs that the female lays for them, which makes the species one of the most difficult poison frogs to breed. It is critically endangered, and the delimitation of the species was changed in 2018.

Família: Dendrobatidae Nome anterior: *Dendrobates histrionicus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**3-4 cm** comprimento focinho-cloaca
Longevidade	**9 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Especialista**

CUIDADOS

Terrário mín.	**50 × 50 × 50 cm** C × L × A para 2 adultos
Temperatura, dia	**24-26 °C**
Temperatura, noite	**18-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**12 meses**
Ovos por postura	**4-20**
Eclosão	**10 dias**
Fase larval	**90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ovos não fertilizados da fêmea**
Cuidado parental	**Vigia os ovos, carrega os girinos e alimenta os girinos com ovos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo crítico (CR)**

Resumo dos cuidados

O TERRÁRIO	A pair needs a rainforest vivarium of at least 50 × 50 × 50 cm, and larger is better. *Oophaga* species can be territorial, so keep only one pair unless the vivarium is very large.
TEMPERATURA	The daytime temperature should be 24-26 °C, and at night it is lowered by 5-6 degrees, to 18-21 °C. The temperature only needs to vary by 1-2 degrees over the year, and there is no winter rest.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-80 % in the middle of the day and up to 100 % morning and evening. This is easiest to achieve with a rain system and a fogging system on timers.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish with about 1 cm of water and a ramp of stones or moss is enough for the adults. Use reverse-osmosis water, rain water or dechlorinated tap water, and change the water often.
ILUMINAÇÃO E UVB	The lighting is first and foremost for the bromeliads and the other plants. Provide about 12 hours of light a day with LED or fluorescent tubes that do not give off much heat.
ALIMENTAÇÃO	*Oophaga histrionica* eats small arthropods, in the wild especially ants, termites, mites and small beetles.
COMPORTAMENTO E MANUSEIO	*Oophaga histrionica* is diurnal. The male calls from a low perch, and males defend their territories. Therefore keep only one pair in the enclosure. Do not handle the frogs.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia.