herpy

SAPOS-FLECHA

Lehmann's Poison Frog

Oophaga lehmanni

Oophaga lehmanni is a rare, diurnal poison frog from a small area in western Colombia, black or dark brown with broad bands of red, orange or yellow. It is critically endangered in the wild, and only captive-bred animals belong in an enclosure. The species requires an experienced keeper: it does not tolerate heat, and the tadpoles live exclusively on unfertilised eggs from the female.

Família: Dendrobatidae Nome anterior: *Dendrobates lehmanni*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**3-4 cm** comprimento focinho-cloaca
Longevidade	**4-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical e floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**60 × 60 × 60 cm** C × L × A para 2 adultos
Temperatura, dia	**22-24 °C**
Temperatura, noite	**17-19 °C**
Umidade	**70-100%**
Água	**Sem recipiente de água**
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Em folhas sobre a água**
Ovos por postura	**3-10**
Eclosão	**14-28 dias**
Fase larval	**60-90 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ovos não fertilizados da fêmea**
Cuidado parental	**Vigia os ovos, carrega os girinos e alimenta os girinos com ovos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo crítico (CR)**

Resumo dos cuidados

O TERRÁRIO	A pair of *O. lehmanni* needs a vivarium of at least 60 × 60 × 60 cm (length × width × height), and larger is better, especially for breeding.
TEMPERATURA	*O. lehmanni* comes from cool mountain rainforest and is more sensitive to heat than most poison frogs. It does best at 22-24 °C during the day, and 24-25 °C should not be exceeded, not even briefly.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %: high in the morning and evening, preferably with fog so that it reaches 100 %, and a little lower in the middle of the day, down to about 70 %.
ÁGUA E QUALIDADE DA ÁGUA	The frogs do not need a water bowl, because the humidity covers their needs.
ILUMINAÇÃO E UVB	The frog is diurnal and lives in shade under the forest canopy. There is no documentation that the species needs UVB when the food is dusted with vitamin D3, but weak UVB lighting (Ferguson zone 1) can be used without harm.
ALIMENTAÇÃO	*O. lehmanni* eats only live, small arthropods. The staple food is flightless fruit flies, and the species is large enough to take the larger *Drosophila hydei* as well.
COMPORTAMENTO E MANUSEIO	*O. lehmanni* is diurnal and calm, but the males are territorial and call from fixed spots. Keep the species in pairs.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia.