herpy

SAPOS-FLECHA

Strawberry Poison Frog

Oophaga pumilio

Oophaga pumilio is a small, diurnal poison frog of 17-24 mm from the rainforests of Central America, known for its many colour forms and an unusual form of parental care: the female feeds each tadpole with unfertilised eggs. It is fascinating to watch, but requires experience, small feeder insects and patience.

Família: Dendrobatidae Nome anterior: *Dendrobates pumilio*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**17-24 mm** comprimento focinho-cloaca
Longevidade	**10-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**10 meses**
Ovos por postura	**3-5**
Eclosão	**10-14 dias**
Fase larval	**42-56 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ovos não fertilizados da fêmea**
Cuidado parental	**Vigia os ovos, carrega os girinos e alimenta os girinos com ovos**

CUIDADOS

Terrário mín.	**45 × 45 × 45 cm** C × L × A para 4 adultos
Temperatura, dia	**22-26 °C**
Temperatura, noite	**19-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	A pair or a small group of up to four animals needs a vivarium of at least 45 × 45 × 45 cm (length × width × height). More animals require more floor area.
TEMPERATURA	By day the temperature should be 22-26 °C, preferably about 24 °C, and at night 19-21 °C. Temperatures below 18 °C or above 32 °C for any length of time can be fatal.
UMIDADE E NEBULIZAÇÃO	Humidity should be high, 70-100 %. Aim for 70-80 % during the day and up to 100 % at night, for example by misting four times a day with an automatic system. A timer-controlled rain system is therefore a great advantage.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish is enough, but the water must be no more than about ¾ of the frog's height, in practice under 1 cm, because the species swims poorly and can drown.
ILUMINAÇÃO E UVB	Give 12 hours of light a day, so that the plants thrive. An LED plant lamp that does not give off much heat is a good choice.
ALIMENTAÇÃO	*Oophaga pumilio* needs very small feeder insects, at most about 2 mm. Small, flightless *Drosophila melanogaster* and springtails are the staple, supplemented with aphids, termites and other small feeders.
COMPORTAMENTO E MANUSEIO	*Oophaga pumilio* is diurnal, bold and inquisitive and moves about a great deal in the leaf litter. The males defend territories with a buzzing, cricket-like call and may fight other males.
ÁREA DE OCORRÊNCIA NATURAL	Nicarágua, Costa Rica e Panamá.