herpy

SAPOS-FLECHA

Vicente's Poison Frog

Oophaga vicentei

Oophaga vicentei is a small, brightly coloured and diurnal poison dart frog of up to 24 mm, found only in central Panama. The colours vary from yellow and red to blue, grey and green, often with a black reticulated pattern. The species is endangered in the wild, and little is known about its husbandry.

Família: Dendrobatidae Nome anterior: *Dendrobates vicentei*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**12-24 mm** comprimento focinho-cloaca
Longevidade	**10-15 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Nas axilas das folhas (por exemplo, bromélias)**
Ovos por postura	**1-12**
Eclosão	**10-14 dias**
Fase larval	**42-56 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ovos não fertilizados da fêmea**
Cuidado parental	**Vigia os ovos, carrega os girinos e alimenta os girinos com ovos**

CUIDADOS

Terrário mín.	**45 × 45 × 60 cm** C × L × A para 2 adultos
Temperatura, dia	**22-26 °C**
Temperatura, noite	**19-21 °C**
Umidade	**80-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo (EN)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the husbandry of *Oophaga vicentei*.
TEMPERATURA	The daytime temperature should be 22-26 °C and the night 19-21 °C. The frogs tolerate heat poorly, so avoid longer periods above 28 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 % and never below 70 %. Mist the enclosure thoroughly once or twice a day with water free of lime and chlorine (rainwater or reverse-osmosis water), so that leaves and bromeliads are wet.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water bowl with 1-2 cm of water is enough, because the frogs are poor swimmers. Keep it clean, and change the water every day or every other day.
ILUMINAÇÃO E UVB	UVB is hardly necessary, but a little UVB (Ferguson zone 1) does no harm and is optional. Plant lighting for 10-12 hours a day keeps the plants alive, and the night must be dark. The lamps must not heat the enclosure above 26 °C.
ALIMENTAÇÃO	The species needs very small feeder insects. The staple is small, flightless fruit flies (*Drosophila melanogaster*) and springtails, supplemented with aphids, small termites and small isopods.
COMPORTAMENTO E MANUSEIO	*Oophaga vicentei* is diurnal. The male calls from a spot high up and finds the breeding site. Females may eat other females' eggs, so a pair is the safest. The frogs should not be handled, because the skin is thin and sensitive.
ÁREA DE OCORRÊNCIA NATURAL	Panamá.