herpy

SAPOS-FLECHA

Kokoe Poison Frog

Phyllobates aurotaenia

Phyllobates aurotaenia is a diurnal poison frog of 3-4 cm from the Pacific coast of Colombia, black with two green to golden stripes along the back. Wild animals carry a deadly skin poison, but captive-bred animals are non-poisonous. The species is bold, calls loudly and is easy to breed.

Família: Dendrobatidae Nome anterior: *Dendrobates aurotaenia*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**3-4 cm** comprimento focinho-cloaca
Longevidade	**10-20 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Fortes**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**8-10 meses**
Ovos por postura	**8-28**
Fase larval	**60-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa**
Cuidado parental	**Carrega os girinos**

CUIDADOS

Terrário mín.	**60 × 45 × 45 cm** C × L × A para 4 adultos
Temperatura, dia	**21-24 °C**
Temperatura, noite	**18-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	A pair or a small group of *Phyllobates aurotaenia* needs a rainforest enclosure of at least 60 × 45 × 45 cm for up to 4 adults. The species is active and also uses height, so give it perches and plants to climb in.
TEMPERATURA	*Phyllobates aurotaenia* does best when cool. During the day it should be 21-24 °C, and at night the temperature is lowered to 18-21 °C. The temperature must not exceed 27 °C, because the genus tolerates heat poorly.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %. The species tolerates down to 50 % briefly if it has access to water, but low humidity without water can quickly become fatal.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish with 1-2 cm of water and a stone or a sloping edge, so the frogs can easily get up, is enough.
ILUMINAÇÃO E UVB	The light is first and foremost for the plants. Fluorescent tubes or LED light of about 6,500 K give natural colours and good plant growth, and about 12 hours of light a day is usual.
ALIMENTAÇÃO	*Phyllobates aurotaenia* eats small arthropods. Juveniles take wingless *Drosophila melanogaster*, while adults readily eat larger feeder insects such as *Drosophila hydei* and very small crickets.
COMPORTAMENTO E MANUSEIO	*Phyllobates aurotaenia* is diurnal and bold, but shy when young.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia.

O guia completo com mapa e atualizações: herpy.org/en/poison-dart-frogs/phyllobates-aurotaenia