herpy

SAPOS-FLECHA

Black-legged Poison Dart Frog

Phyllobates bicolor

Phyllobates bicolor is a yellow to orange poison frog of 3-5 cm from the rainforest of western Colombia, often with black or greenish legs. In the wild it is one of the most poisonous animals known, but captive-bred animals are non-poisonous. It is bold, curious and almost always on view, and it breeds readily in the enclosure.

Família: Dendrobatidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**3-5 cm** comprimento focinho-cloaca
Peso	**4,5 g**
Longevidade	**6-20 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Fortes**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**50 × 50 × 50 cm** C × L × A para 2 adultos
Temperatura, dia	**21-25 °C**
Temperatura, noite	**18-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Insetos e pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**12 meses**
Ovos por postura	**4-30**
Eclosão	**13-14 dias** a 21-23 °C
Fase larval	**21-60 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo (EN)**

Resumo dos cuidados

O TERRÁRIO	A pair of *Phyllobates bicolor* needs a rainforest enclosure of at least 50 × 50 × 50 cm. Allow about 40 litres per adult. A group of up to nine animals needs at least 70 × 70 × 70 cm.
TEMPERATURA	During the day it should be 21-25 °C, and at night the temperature is lowered by 3-4 degrees, to 18-21 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-80 % in the middle of the day and up to 100 % morning and evening. A rain system and a fogging system on timers make this easiest, and the fog clearly makes the frogs call and move about.
ÁGUA E QUALIDADE DA ÁGUA	The species appreciates a water surface in the enclosure.
ILUMINAÇÃO E UVB	The lighting is first and foremost for the plants. Fluorescent tubes or LED for about 12 hours a day are enough.
ALIMENTAÇÃO	*Phyllobates bicolor* prefers small feeder insects of no more than about 3 mm.
COMPORTAMENTO E MANUSEIO	*Phyllobates bicolor* is diurnal, very active and curious and almost always on view. The males call loudly with trilling, buzzing tones, sometimes for several days at a time.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia.