herpy

SAPOS-FLECHA

Lovely Poison Frog

Phyllobates lugubris

Phyllobates lugubris is a small, black poison frog of 19-24 mm with two yellow, golden or orange stripes along the back. It lives on the forest floor in the lowland rainforest on the Caribbean side of Central America, from Nicaragua to Panama. It is occasionally seen in the trade, but little is known about its care.

Família: Dendrobatidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**19-24 mm** comprimento focinho-cloaca
Longevidade	**10-20 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**7-21**
Fase larval	**45-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

CUIDADOS

Terrário mín.	**60 × 45 × 45 cm** C × L × A para 3 adultos
Temperatura, dia	**21-25 °C**
Temperatura, noite	**18-21 °C**
Umidade	**80-100%**
Água	**Recipiente de água** profundidade 1-3 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little is known about the care of *Phyllobates lugubris*.
TEMPERATURA	The daytime temperature should be 21-25 °C and the night 18-21 °C, as for *Phyllobates vittatus*. Poison frogs are very sensitive to heat: above 27 °C should be avoided, and several hours at around 29 °C can be fatal.
UMIDADE E NEBULIZAÇÃO	Humidity should be 80-100 %, and the soil and leaf litter must always be moist. Mist the enclosure thoroughly 1-2 times a day with rainwater, reverse-osmosis water or dechlorinated water, or use a rain or mist system.
ÁGUA E QUALIDADE DA ÁGUA	Adult frogs do not drink, but absorb water through the skin, and a shallow water bowl with 1-3 cm of water is enough. *Phyllobates vittatus* appreciates a small water feature, and in the wild the tadpoles are carried to streams.
ILUMINAÇÃO E UVB	The species lives on the forest floor, where the light is subdued, and UVB is not documented as necessary. A weak UVB lamp in Ferguson zone 1 is optional; if used, always provide shade.
ALIMENTAÇÃO	In the wild *Phyllobates lugubris* eats mostly ants and mites. In the enclosure, small arthropods are given: wingless fruit flies (*Drosophila melanogaster*), springtails, small isopods and newly hatched crickets.
COMPORTAMENTO E MANUSEIO	The species is diurnal and lives hidden among leaf litter and plants. Males call from hidden spots and defend their territory against other males, so several males in the same enclosure can cause stress.
ÁREA DE OCORRÊNCIA NATURAL	Nicarágua, Costa Rica e Panamá.