herpy

SAPOS-FLECHA

Golden Poison Frog

Phyllobates terribilis

Phyllobates terribilis is the largest of the poison frogs, 4-5 cm, and in the wild one of the most toxic animals known. It lives only in a small patch of rainforest on Colombia's Pacific coast. Captive-bred animals are practically non-toxic, bold and always on show, and the species has become one of the most commonly kept poison frogs in the hobby.

Família: Dendrobatidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**4-5 cm** comprimento focinho-cloaca
Longevidade	**8-15 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Fortes**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**60 × 50 × 50 cm** C × L × A para 2 adultos
Temperatura, dia	**21-25 °C**
Temperatura, noite	**18-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Insetos e pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**16-20 meses**
Ovos por postura	**10-30**
Eclosão	**10-25 dias** a 22-24 °C
Fase larval	**30-60 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo (EN)**

Resumo dos cuidados

O TERRÁRIO	A pair of *Phyllobates terribilis* needs an enclosure of at least 60 × 50 × 50 cm. A group of 4-6 animals needs at least 90 × 50 × 50 cm, and more animals require more floor area.
TEMPERATURA	During the day it should be 21-25 °C, preferably a couple of degrees warmer at the top of the enclosure, and at night 18-21 °C, which corresponds to room temperature. The temperature must not exceed 27 °C for any length of time.
UMIDADE E NEBULIZAÇÃO	Humidity should be high, 70-100 %, but the floor must not become boggy. A rain system is the easiest, for example with three mistings a day of 20-60 seconds, depending on the nozzles.
ÁGUA E QUALIDADE DA ÁGUA	Provide one or more small water dishes with about 2 cm of water, for example saucers or halved Brazil nut shells, which the animals can easily climb out of.
ILUMINAÇÃO E UVB	The light is first and foremost for the plants. Fluorescent tubes or LED bars for about 12 hours a day are enough, and the day length may vary slightly over the year.
ALIMENTAÇÃO	*Phyllobates terribilis* takes larger prey than most poison dart frogs. In the wild ants are the main food, but it takes many kinds of small arthropods.
COMPORTAMENTO E MANUSEIO	*Phyllobates terribilis* is diurnal, bold and inquisitive and is out on show most of the day with a short break in the middle of the day. The males call with a soft, bird-like trill, especially after misting.
ÁREA DE OCORRÊNCIA NATURAL	Colômbia.