herpy

SAPOS-FLECHA

Golfodulcean Poison Frog

Phyllobates vittatus

Phyllobates vittatus is a black poison frog with two bright orange or yellow stripes down the back and blue marbled legs. It occurs only in the lowland forest around the Golfo Dulce in south-western Costa Rica. In the terrarium it is diurnal, vocal and a prolific breeder, and the male carries the tadpoles to water himself.

Família: Dendrobatidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**3-4 cm** comprimento focinho-cloaca
Longevidade	**10-20 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Maturidade sexual	**8-10 meses**
Ovos por postura	**7-21**
Fase larval	**45-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

CUIDADOS

Terrário mín.	**60 × 45 × 45 cm** C × L × A para 4 adultos
Temperatura, dia	**21-25 °C**
Temperatura, noite	**18-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional** Zona de Ferguson 1
Alimentação	**Insetos e pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	Up to four *Phyllobates vittatus* need an enclosure of at least 60 × 45 × 45 cm. A group of 5-6 animals needs more space, e.g. 100 × 60 × 60 cm, which also gives room both to move on the ground and to climb.
TEMPERATURA	During the day *Phyllobates vittatus* needs 21-25 °C and at night 18-21 °C, which most living rooms provide.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %. The species tolerates down to 50 % for short periods if it has access to water, but low humidity without water can quickly become fatal.
ÁGUA E QUALIDADE DA ÁGUA	Provide a shallow water dish or a small water feature with 1-2 cm of water and a stone or a piece of bark so the animals can climb out easily. The frogs moisten themselves in the water, and the male may use it to deposit tadpoles.
ILUMINAÇÃO E UVB	The lighting is for the plants. LED bars or fluorescent tubes on a timer for 10-12 hours a day are enough, and the night must be dark. Provide shade, and never place the enclosure in direct sun.
ALIMENTAÇÃO	*Phyllobates vittatus* is unusual in that adults take larger prey than most poison frogs.
COMPORTAMENTO E MANUSEIO	*Phyllobates vittatus* is diurnal, and the males have a high, trilling call. The species can be kept in pairs or groups.
ÁREA DE OCORRÊNCIA NATURAL	Costa Rica.