herpy

SAPOS-FLECHA

Amazonian Poison Frog

Ranitomeya amazonica

Ranitomeya amazonica is a tiny, diurnal poison frog of 16-19 mm from the Amazon around Iquitos in Peru and Leticia in Colombia. It is black with red, orange or yellow stripes and blue-grey marbling on the legs, breeds in bromeliads and is regarded as one of the easiest small Ranitomeya to keep and breed.

Família: Dendrobatidae Nome anterior: *Dendrobates amazonicus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**16-19 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Nas axilas das folhas (por exemplo, bromélias)**
Maturidade sexual	**5-8 meses**
Ovos por postura	**2-8**
Eclosão	**10-14 dias**
Fase larval	**60-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa e outros girinos**
Cuidado parental	**Carrega os girinos**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 3 adultos
Temperatura, dia	**21-26 °C**
Temperatura, noite	**18-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	A pair or trio of *Ranitomeya amazonica* needs an enclosure of at least 40 × 40 × 40 cm. If there are to be more frogs, allow about 20 litres of extra space per animal, and choose a tall rather than a low enclosure.
TEMPERATURA	Keep the enclosure at 21-26 °C by day and 18-21 °C at night. The frogs are most active and breed best at the warm end, at 24-26 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %, preferably 80-100 %. Mist every day, preferably morning and evening, so that the humidity is close to 100 % after misting and falls to about 70-75 % in the middle of the day.
ÁGUA E QUALIDADE DA ÁGUA	*Ranitomeya amazonica* swims poorly, so the water must be shallow and easy to climb out of.
ILUMINAÇÃO E UVB	The frogs are diurnal and live in the shade of the forest. The lighting is first and foremost for the plants: give 12-14 hours of LED plant lighting a day. *Ranitomeya amazonica* has no known need for UVB, so UVB is optional.
ALIMENTAÇÃO	*Ranitomeya amazonica* is so small that it can only take small feeder insects. Juveniles and adults take wingless fruit flies (*Drosophila melanogaster*), and large adults can also take *D. hydei* and newly hatched crickets.
COMPORTAMENTO E MANUSEIO	*Ranitomeya amazonica* is diurnal, but often shy, and young animals in particular hide a lot. In a densely planted enclosure with many dead leaves the frogs come out more.
ÁREA DE OCORRÊNCIA NATURAL	Peru e Colômbia.