herpy

SAPOS-FLECHA

Blessed Poison Frog

Ranitomeya benedicta

Ranitomeya benedicta is a small, diurnal poison frog of 16-20 mm from the lowland forest of northern Peru. The bright red head and the black body with a blue net pattern make it one of the most colourful in the genus. Until 2008 it was regarded as a subspecies of *Ranitomeya fantastica*. In the wild it is vulnerable, but captive-bred animals breed willingly in a well-furnished enclosure.

Família: Dendrobatidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**16-20 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**21-26 °C**
Temperatura, noite	**18-22 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**2-6**
Eclosão	**10-14 dias**
Fase larval	**60-114 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	A pair of *R. benedicta* can be kept in an enclosure of 40 × 40 × 40 cm. Two pairs need a horizontal enclosure of about 75 litres, and bigger is always better, because a large enclosure keeps the climate more stable.
TEMPERATURA	Keep the enclosure at 21-26 °C by day and 18-22 °C at night. The species thrives best at about 21-24 °C, and the temperature must not exceed 27 °C for long periods.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %. A good rhythm is almost 100 % morning and evening and 70-75 % in the middle of the day, so the enclosure dries out a little between mistings.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish or a bromeliad with water in the leaf axils is enough.
ILUMINAÇÃO E UVB	*R. benedicta* is diurnal, and the lighting is first and foremost for the plants. Give 12-14 hours of light a day with fluorescent tubes or LED on a timer, all year round, and darkness at night.
ALIMENTAÇÃO	The staple food is small live arthropods. Animals of all ages take flightless fruit flies (*Drosophila melanogaster*), and adults can also take the larger *D. hydei*.
COMPORTAMENTO E MANUSEIO	*R. benedicta* is diurnal and can be a little shy, especially as a juvenile, but in a densely planted enclosure with hiding places adult animals are seen a lot. The male sits openly and calls.
ÁREA DE OCORRÊNCIA NATURAL	Peru.