herpy

SAPOS-FLECHA

Red-headed Poison Frog

Ranitomeya fantastica

Ranitomeya fantastica is a diurnal poison frog of 20-25 mm from northern Peru and one of the largest in the genus. It is unusually variable, and many forms have a bright orange, red or yellow head above a black body with light markings. The frogs are lightning-fast and often shy, the eggs are hard to find, and the species is best suited to a keeper who already has experience with poison frogs.

Família: Dendrobatidae Nome anterior: *Dendrobates fantasticus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**20-25 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 3 adultos
Temperatura, dia	**21-26 °C**
Temperatura, noite	**18-22 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**3-8**
Eclosão	**10-14 dias**
Fase larval	**60-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Vulnerável (VU)**

Resumo dos cuidados

O TERRÁRIO	A pair or trio of *R. fantastica* can be kept in an enclosure of 40 × 40 × 40 cm. Bigger is better. Many forms are arboreal and use the height.
TEMPERATURA	Keep the enclosure at 21-26 °C by day and 18-22 °C at night. The temperature must not exceed 27 °C for long periods, and above 29 °C can be fatal. Heat is more dangerous than cold.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %. A good rhythm is almost 100 % morning and evening and 70-75 % in the middle of the day, so the enclosure is allowed to dry out a little between mistings.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish or a bromeliad with water in the leaf axils is enough.
ILUMINAÇÃO E UVB	*R. fantastica* is diurnal, and the lighting is first and foremost for the plants. Give 12-14 hours of light a day and darkness at night. It is not documented that the species needs UVB.
ALIMENTAÇÃO	The staple food is small arthropods. Juveniles and adults take wingless fruit flies (*Drosophila melanogaster*), and large adults can also take the larger *D. hydei*.
COMPORTAMENTO E MANUSEIO	*R. fantastica* is diurnal, lightning-fast and often shy, but many animals become bolder when they breed. You see most of them in a densely planted enclosure with hiding places.
ÁREA DE OCORRÊNCIA NATURAL	Peru.