herpy

SAPOS-FLECHA

Red-backed Poison Frog

Ranitomeya reticulata

Ranitomeya reticulata is one of the smallest poison frogs: only 13-17 mm, with a coppery red to orange back and blue-grey legs with a black net pattern. It lives in the lowland forest of the Amazon in north-eastern Peru and eastern Ecuador. In the terrarium it is bold and lively, but demanding, because the eggs are hard to find and the juveniles are hard to raise.

Família: Dendrobatidae Nome anterior: *Dendrobates reticulatus*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**13-17 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**30 × 30 × 45 cm** C × L × A para 2 adultos
Temperatura, dia	**22-26 °C**
Temperatura, noite	**18-22 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**1-8**
Eclosão	**10-14 dias**
Fase larval	**60-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	A pair of *R. reticulata* needs an enclosure of at least 30 × 30 × 45 cm, preferably 40 × 40 × 45 cm or larger. The frogs like to climb, so the height is used.
TEMPERATURA	Keep the enclosure at 22-26 °C during the day and 18-22 °C at night. The temperature must not exceed 27 °C for any length of time, and above 29 °C can be fatal. Heat is always more dangerous than cold.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %. A good rhythm is almost 100 % morning and evening and 70-75 % in the middle of the day, so the enclosure dries out a little between sprayings.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish or a bromeliad with water in the leaf axils is enough.
ILUMINAÇÃO E UVB	*R. reticulata* is diurnal, and the lighting is first and foremost for the plants. Give 12-14 hours of light a day, e.g. LED from 8 to 20 hrs, and darkness at night. It is not documented that the species needs UVB.
ALIMENTAÇÃO	The staple diet is small arthropods. Juveniles and most adults take flightless fruit flies (*Drosophila melanogaster*), and large adults can also take the larger *D. hydei*.
COMPORTAMENTO E MANUSEIO	*R. reticulata* is lightning fast and bold and is often seen out in the open, as if it has no idea how small it is. Males call from fixed spots with a high, short buzz.
ÁREA DE OCORRÊNCIA NATURAL	Peru e Equador.

O guia completo com mapa e atualizações: herpy.org/en/poison-dart-frogs/ranitomeya-reticulata