herpy

SAPOS-FLECHA

Sira Poison Frog

Ranitomeya sirensis

Ranitomeya sirensis is a small, diurnal poison frog of 15-25 mm from eastern Peru, northern Bolivia and the far west of Brazil. The species includes the forms that were formerly sold as "lamasi" and "biolat", and most of them are among the easiest poison frogs to breed. The frogs thrive in groups when there is enough space.

Família: Dendrobatidae Nome anterior: *Ranitomeya lamasi; Dendrobates lamasi; Dendrobates sirensis*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**15-25 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical e floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**45 × 45 × 45 cm** C × L × A para 3 adultos
Temperatura, dia	**21-25 °C**
Temperatura, noite	**18-21 °C**
Umidade	**80-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Nas axilas das folhas (por exemplo, bromélias)**
Maturidade sexual	**5-8 meses**
Ovos por postura	**1-8**
Eclosão	**10-14 dias**
Fase larval	**58-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO
A pair or trio of *R. sirensis* needs an enclosure of at least 45 × 45 × 45 cm. The species benefits from a little extra floor space and dense plant cover, and the large highland form thrives best in a larger enclosure.

TEMPERATURA
Keep the enclosure at 21-25 °C during the day and 18-21 °C at night. The temperature must not exceed 27 °C for any length of time, and above 29 °C is dangerous.

UMIDADE E NEBULIZAÇÃO
Humidity should be 80-100 %. The species tolerates down to 50 % only briefly, and only if it has access to water; low humidity without water can quickly become fatal.

ÁGUA E QUALIDADE DA ÁGUA
A shallow water dish or a bromeliad with water in the leaf axils is enough. Keep the water 1-2 cm deep, so the smallest frog can sit in it with its head and upper body above the water.

ILUMINAÇÃO E UVB
R. sirensis is diurnal, and the lighting is first and foremost for the plants. Give 12-14 hours of light a day with fluorescent tubes or LED and darkness at night. It is not documented that the species needs UVB.

ALIMENTAÇÃO
The staple diet is small arthropods: wingless fruit flies (*Drosophila melanogaster*), springtails and small woodlice. Adults also take the larger *D. hydei* and newly hatched crickets.

COMPORTAMENTO E MANUSEIO
R. sirensis is often shy and likes to hide, so a densely planted enclosure with many hiding places gives the most to look at. The male calls to lure a female to an egg-laying site.

ÁREA DE OCORRÊNCIA NATURAL
Bolívia, Brasil e Peru.