herpy

SAPOS-FLECHA

Summers' Poison Dart Frog

Ranitomeya summersi

Ranitomeya summersi is a small, diurnal poison frog of 16-21 mm from the Huallaga Valley in Peru, orange or copper-coloured with black marbling and a black mask over the eyes. The species is threatened in the wild, the eggs are hard to find, and the frogs are quick and shy, so it is best suited to a keeper with experience of poison frogs.

Família: Dendrobatidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**16-21 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**30 × 30 × 45 cm** C × L × A para 3 adultos
Temperatura, dia	**21-26 °C**
Temperatura, noite	**18-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**No chão**
Ovos por postura	**4-9**
Eclosão	**10-14 dias**
Fase larval	**60-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Vigia os ovos e carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Em perigo (EN)**

Resumo dos cuidados

O TERRÁRIO	A pair or a small trio of *R. summersi* can be kept in an enclosure of at least 30 × 30 × 45 cm (length × width × height), and bigger is better.
TEMPERATURA	Keep the enclosure at 21-26 °C during the day and 18-21 °C at night. The temperature must not exceed 27 °C for any length of time, and above 29 °C is life-threatening. The species tolerates heat worse than cold.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %. A good rhythm is close to 100 % morning and evening and 70-75 % in the middle of the day, so the enclosure can dry out a little between mistings.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish or a bromeliad with water in the leaf axils is enough.
ILUMINAÇÃO E UVB	*R. summersi* is diurnal, and the light is first and foremost for the plants. Give 12-14 hours of light a day and darkness at night. It is not documented that the species needs UVB.
ALIMENTAÇÃO	The staple food is small live arthropods. Juveniles and young animals take wingless fruit flies (*Drosophila melanogaster*), and large adults can also take the larger *D. hydei*.
COMPORTAMENTO E MANUSEIO	*R. summersi* is diurnal, quick and often shy, but can become bolder once it has got used to the enclosure. They are seen most in a densely planted enclosure with hiding places.
ÁREA DE OCORRÊNCIA NATURAL	Peru.