herpy

SAPOS-FLECHA

Red Uakari Poison Frog

Ranitomeya uakarii

Ranitomeya uakarii is a very small, diurnal poison frog of 13-16 mm from the Amazon basin. It is black with red to orange stripes, yellow flanks and blue legs, and it is rare in the hobby. The care advice is based on experience with the genus, and the species suits an experienced keeper with patience and a steady supply of tiny feeder insects.

Família: Dendrobatidae Nome anterior: *Dendrobates uakarii*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**13-16 mm** comprimento focinho-cloaca
Longevidade	**5-6 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**45 × 45 × 45 cm** C × L × A para 3 adultos
Temperatura, dia	**21-25 °C**
Temperatura, noite	**18-21 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Nas axilas das folhas (por exemplo, bromélias)**
Ovos por postura	**2-7**
Eclosão	**10-14 dias**
Fase larval	**60-100 dias** da eclosão até a metamorfose
Alimentação das larvas	**Algas/plantas e ração completa**
Cuidado parental	**Carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	A pair or trio of *Ranitomeya uakarii* needs an enclosure of at least 45 × 45 × 45 cm. The size is borrowed from the genus. If you use juveniles, the safest approach is to start with at least four, so that both sexes are included.
TEMPERATURA	Keep the enclosure at 21-25 °C during the day and 18-21 °C at night. The figures are borrowed from the genus. Extra heat is rarely necessary in a heated living room.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %. Mist daily, so that humidity is close to 100 % morning and evening and 70-75 % in the middle of the day. This lets the terrarium dry out a little between sprayings.
ÁGUA E QUALIDADE DA ÁGUA	A shallow water dish or a bromeliad with water in the leaf axils is enough. Keep the water 1-2 cm deep, so that the small frog can sit in it with its head and upper body above the water; the depth is borrowed from the genus.
ILUMINAÇÃO E UVB	*Ranitomeya uakarii* is diurnal, and the lighting is first and foremost for the plants. Provide 12-14 hours of light a day with fluorescent tubes or LED and darkness at night. It is not documented that the species needs UVB.
ALIMENTAÇÃO	The staple diet is small arthropods: wingless fruit flies, springtails, small white woodlice and newly hatched crickets.
COMPORTAMENTO E MANUSEIO	*Ranitomeya uakarii* is diurnal. In a densely planted terrarium with many hiding places you typically see more of the frogs than in a barren terrarium, but the small frogs are often shy and hide when you approach the glass.
ÁREA DE OCORRÊNCIA NATURAL	Brasil, Colômbia e Peru.