herpy

SAPOS-FLECHA

Brazilian Poison Frog

Ranitomeya vanzolinii

Ranitomeya vanzolinii is a small, diurnal poison frog of 17-19 mm from the western Amazon. It lives in the trees and rears its young in water that collects in bromeliads. The female feeds the tadpoles with unfertilised eggs.

Família: Dendrobatidae Nome anterior: *Dendrobates vanzolinii*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**17-19 mm** comprimento focinho-cloaca
Longevidade	**5-6 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola**
Habitat	**Floresta tropical e floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Nas axilas das folhas (por exemplo, bromélias)**
Ovos por postura	**1-6**
Eclosão	**10-14 dias**
Fase larval	**60-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa e ovos não fertilizados da fêmea**
Cuidado parental	**Carrega os girinos e alimenta os girinos com ovos**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**23-26 °C**
Temperatura, noite	**19-23 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO
A pair of *Ranitomeya vanzolinii* needs an enclosure of at least 40 × 40 × 40 cm, preferably 50 × 50 × 50 cm. The frog is arboreal and uses the height, so a tall enclosure with many vertical surfaces is better than a low, wide one.

TEMPERATURA
The daytime temperature should be 23-26 °C, and at night it is lowered by 3-4 °C to 19-23 °C. The variation over the year should be small, 1-2 °C. A basking spot is not necessary, and the species does not hibernate.

UMIDADE E NEBULIZAÇÃO
Humidity should be 70-100 %. Morning and evening it should be close to 100 %, and at midday it may fall to 70-75 %. The terrarium is misted daily, preferably morning and evening, with reverse-osmosis or rain water.

ÁGUA E QUALIDADE DA ÁGUA
A water dish is not necessary for the adults, but is an advantage, and it can be shallow with 1-2 cm of water.

ILUMINAÇÃO E UVB
The lighting requirement for *Ranitomeya vanzolinii* is modest. The frog needs enough light to see its food, and the plants need more. LED lighting for plants or a fluorescent tube, switched on 12-14 hours a day, is suitable.

ALIMENTAÇÃO
Ranitomeya vanzolinii feeds on small, live arthropods: fruit flies, springtails, small tropical isopods and newly hatched crickets. Small animals require small feeder insects.

COMPORTAMENTO E MANUSEIO
Ranitomeya vanzolinii is diurnal, but shy. In a densely planted terrarium with many hiding places it shows itself more often than in a sparsely furnished one.

ÁREA DE OCORRÊNCIA NATURAL
Brasil e Peru.