herpy

SAPOS-FLECHA

Variable Poison Frog

Ranitomeya variabilis

Ranitomeya variabilis is a small, diurnal poison frog of 15-20 mm from the eastern slopes of the Andes and the Amazon. It has many local forms in yellow, green and blue with black markings, breeds in bromeliads and is known for the male carrying the tadpoles around between the plant's pools of water. A good first species among the small poison frogs.

Família: Dendrobatidae Nome anterior: *Dendrobates variabilis*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**15-20 mm** comprimento focinho-cloaca
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola e terrestre**
Habitat	**Floresta tropical e floresta de montanha**
Toxinas na pele na natureza	**Leves**
Toxinas na pele em cativeiro	**Nenhuma** Animais nascidos em cativeiro. Animais capturados na natureza podem manter as toxinas por muito tempo
Dificuldade	**Intermediário**

CUIDADOS

Terrário mín.	**40 × 40 × 40 cm** C × L × A para 2 adultos
Temperatura, dia	**21-26 °C**
Temperatura, noite	**18-22 °C**
Umidade	**70-100%**
Água	**Recipiente de água** profundidade 1-2 cm
UVB	**Opcional**
Alimentação	**Pequenos invertebrados (colêmbolos, drosófilas)**
Mantido	**Em casal ou em grupo**

REPRODUÇÃO

Reprodução	**Ovíparo**
Desenvolvimento	**Girinos carregados até a água**
Local de postura	**Nas axilas das folhas (por exemplo, bromélias)**
Ovos por postura	**1-10**
Eclosão	**10-14 dias**
Fase larval	**60-80 dias** da eclosão até a metamorfose
Alimentação das larvas	**Ração completa e outros girinos**
Cuidado parental	**Carrega os girinos**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	A pair of *R. variabilis* needs an enclosure of at least 40 × 40 × 40 cm, and the same size is enough for a trio. Each extra frog needs approx. 4 litres more space.
TEMPERATURA	Keep the enclosure at 21-26 °C during the day and 18-22 °C at night. For short periods the species tolerates up to 27 °C, but the temperature must never exceed 29 °C. Eggs and tadpoles can be kept at a constant 24 °C.
UMIDADE E NEBULIZAÇÃO	Humidity should be 70-100 %. Aim for almost 100 % morning and evening and 70-75 % in the middle of the day. The species can cope briefly at 50 % if it has clean water to sit in, but this is an emergency measure.
ÁGUA E QUALIDADE DA ÁGUA	Give the frogs a shallow water dish or a small water feature, and keep the water so shallow that the smallest frog can sit with its head and upper body above the water; the species swims poorly.
ILUMINAÇÃO E UVB	The frogs are diurnal and live in the shade of the forest. The lighting is first and foremost for the plants: LED light for 12-14 hours a day is suitable.
ALIMENTAÇÃO	*R. variabilis* is so small that it can only take small feeder insects: springtails, wingless fruit flies (*Drosophila melanogaster*) and small woodlice. Large adults can also take *D. hydei*.
COMPORTAMENTO E MANUSEIO	*R. variabilis* is diurnal and peaceful. The males call from elevated spots, and in the wild the species has a free mating system without fixed pairs and without defended territories.
ÁREA DE OCORRÊNCIA NATURAL	Peru, Equador, Colômbia, Brasil e Guiana Francesa.