herpy

COBRAS

Brown House Snake

Boaedon capensis

Boaedon capensis is a slender, nocturnal constrictor from southern Africa, where it lives in scrub, grassland and around human dwellings and hunts rodents and lizards. In the enclosure it is hardy and simple to keep, and adults usually become calm in the hand, but the species is a skilled escape artist, and juveniles are fast.

Família: Lamprophiidae Nome anterior: *Lamprophis capensis; Boaedon capense*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**60-120 cm** comprimento total com a cauda
Longevidade	**10-20 anos**
Ativo	**À noite**
Estilo de vida	**Terrestre**
Habitat	**Semideserto e savana**
Dificuldade	**Iniciante**

CUIDADOS

Terrário mín.	**90 × 45 × 40 cm** C × L × A para 1 adulto
Temperatura, dia	**24-28 °C**
Ponto quente	**30-32 °C**
Temperatura, noite	**20-24 °C**
Umidade	**40-60%**
Água	**Recipiente de água**
UVB	**Opcional**
Alimentação	**Vertebrados**
Mantido	**Sozinho**

REPRODUÇÃO

Reprodução	**Ovíparo**
Tamanho da postura	**5-16 ovos**
Posturas por ano	**1-4**
Incubação	**55-65 dias** a 29-30 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	An adult *B. capensis* needs an enclosure of at least 90 × 45 × 40 cm. Larger is always better, and large females do well in 100 × 50 × 40 cm.
TEMPERATURA	The enclosure must have a clear temperature gradient. During the day it is 24-28 °C, with about 24 °C at the cool end. The warmest spot, where the snake can lie and warm itself, should be 30-32 °C.
UMIDADE E ÁGUA	Humidity should be 40-60%. In most rooms this is achieved with a water bowl, and it is rarely necessary to mist.
ILUMINAÇÃO E UVB	*B. capensis* is nocturnal and does not need bright light. A fixed light rhythm of about 12 hours of light and 12 hours of dark all year round gives a clear daily rhythm. UVB is not necessary for animals that are fed whole prey.
ALIMENTAÇÃO	*Boaedon capensis* kills its prey by constriction, has a strong feeding response and usually eats readily. It is fed whole, thawed mice and, for large females, small rats, never live prey.
COMPORTAMENTO E MANUSEIO	Juveniles and subadults can be fast, nippy and restless, and they can spring out of the enclosure like a released spring when the lid is opened. Open the lid carefully over a safe spot.
ÁREA DE OCORRÊNCIA NATURAL	África do Sul, Lesoto, Essuatíni, Botsuana, Zimbábue, Moçambique, Zâmbia e Malaui.

O guia completo com mapa e atualizações: herpy.org/en/snakes/boaedon-capensis