herpy

COBRAS

Plains Garter Snake

Thamnophis radix

Thamnophis radix is a slim, striped garter snake from the prairies and open grasslands of central North America. It has a yellow or orange stripe down the back and grows to 40-70 cm. It is very tolerant of cold and gives birth to live young. It is hardy and can be kept on thawed mice, but it escapes through the smallest gaps.

Família: Colubridae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**40-70 cm** comprimento total com a cauda
Longevidade	**5-10 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre**
Habitat	**Área úmida e bosque e prado temperados**
Dificuldade	**Iniciante**

CUIDADOS

Terrário mín.	**100 × 50 × 50 cm** C × L × A para 1 adulto
Temperatura, dia	**23-27 °C**
Ponto quente	**29-32 °C**
Temperatura, noite	**18-24 °C**
Umidade	**35-60%**
Água	**Recipiente de água**
UVB	**Recomendado** Zona de Ferguson 2
Repouso de inverno	**8-15 °C** 8-16 semanas
Alimentação	**Vertebrados e minhocas/caracóis**
Mantido	**Sozinho ou em grupo**

REPRODUÇÃO

Reprodução	**Vivíparo**
Maturidade sexual	**24-36 meses**
Tamanho da ninhada	**5-60 filhotes**
Posturas por ano	**1**
Gestação	**83-102 dias**

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Não listado**
Anexo da UE	**Não listado**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	Little species-specific knowledge exists about keeping *Thamnophis radix*.
TEMPERATURA	There must be a clear temperature gradient. The basking spot should be 29-32 °C measured on the surface, the cool end 23-27 °C and the night 18-24 °C.
UMIDADE E ÁGUA	The air humidity should be 35-60 %, highest at the cool end and at night. The enclosure should be dry, because persistently damp conditions cause blisters on the belly and fungal infections.
ILUMINAÇÃO E UVB	*Thamnophis radix* is diurnal and thrives in strong daylight. UVB is not strictly necessary, but is recommended.
ALIMENTAÇÃO	In the wild *Thamnophis radix* eats earthworms, slugs, frogs, toads, salamanders, fish and the occasional mouse.
COMPORTAMENTO E MANUSEIO	*Thamnophis radix* is quick and alert, but usually becomes calm if handled regularly from a young age. Let new animals settle in for about two weeks, and start handling them only when they are feeding steadily.
ÁREA DE OCORRÊNCIA NATURAL	Canadá e Estados Unidos.