herpy

JABUTIS, CÁGADOS E TARTARUGAS

## Red-Footed Tortoise

# *Chelonoidis carbonarius*

*Chelonoidis carbonarius* is a medium-sized, dark tortoise from the savannas and forest edges of South America with red to orange scales on the head and legs. The shell is typically 30-35 cm long, the animal lives for at least 50 years and is one of the most commonly kept tropical tortoises. It needs high humidity and warmth all year and eats plants, fruit and a little animal food.

Família: Testudinidae  Nome anterior: *Chelonoidis carbonaria; Geochelone carbonaria; Testudo carbonaria*

### DADOS PRINCIPAIS

#### O ANIMAL

| | |
|---|---|
| Tamanho adulto | **30-45 cm**<br>comprimento total com a cauda |
| Longevidade | **50-70 anos** |
| Ativo | **De dia** |
| Estilo de vida | **Terrestre** |
| Habitat | **Savana, floresta seca e floresta tropical** |
| Dificuldade | **Intermediário** |

#### CUIDADOS

| | |
|---|---|
| Terrário mín. | **280 × 140 × 60 cm**<br>C × L × A para 1 adulto |
| Temperatura, dia | **24-29 °C** |
| Ponto quente | **32-35 °C** |
| Temperatura, noite | **20-23 °C** |
| Umidade | **70-90%** |
| Água | **Recipiente de água** |
| UVB | **Essencial**<br>Zona de Ferguson 2 |
| Alimentação | **Insetos, frutas, plantas e minhocas/caracóis** |
| Mantido | **Sozinho ou em grupo** |

#### REPRODUÇÃO

| | |
|---|---|
| Reprodução | **Ovíparo** |
| Tamanho da postura | **2-7 ovos** |
| Incubação | **105-200 dias**<br>a 28-31 °C |

#### PROTEÇÃO E DOCUMENTAÇÃO

| | |
|---|---|
| CITES | **Apêndice II** |
| Anexo da UE | **Anexo B** |
| IUCN | **Não avaliado (NE)** |

## Resumo dos cuidados

| | |
|---|---|
| **O TERRÁRIO** | *Chelonoidis carbonarius* is an active tortoise that uses all the space it is given. It lives on the ground, so floor area is more important than height. |
| **TEMPERATURA** | Indoors there must be a gradient. The air temperature is 24-29 °C by day, and the basking spot under the heat lamp is 32-35 °C, measured on the surface itself. |
| **UMIDADE E ÁGUA** | The species comes from a warm and humid climate and needs high humidity: 70-90 %, with short peaks towards 100 %. |
| **ILUMINAÇÃO E UVB** | *C. carbonarius* is diurnal and needs UVB when it is not in direct sun. Measure a UV index of 2.0-3.0 at the basking spot (Ferguson zone 2). |
| **ALIMENTAÇÃO** | *C. carbonarius* is an omnivore and in the wild eats plants, flowers, fruit, fungi and a little animal food. In captivity a split of about 90 % plant material and 10 % animal food works. |
| **COMPORTAMENTO E MANUSEIO** | The species is inquisitive, active and fairly calm. It gets to know its keeper, readily comes over when there is food and takes food from a hand or tweezers. |
| **ÁREA DE OCORRÊNCIA NATURAL** | Panamá, Colômbia, Venezuela, Guiana, Suriname, Guiana Francesa, Brasil, Peru, Bolívia, Paraguai, Argentina e Trinidad e Tobago. |