herpy

VARANOS

White-throated Monitor

Varanus albigularis

Varanus albigularis is a powerful, diurnal monitor that loves to dig, from savannas and dry woodland in southern and eastern Africa. It grows to 100-200 cm, is inquisitive and can become very tame, but it needs a room rather than an enclosure, a large food bill and many years of careful husbandry.

Família: Varanidae

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**100-200 cm** comprimento total com a cauda
Peso	**3.000-12.000 g**
Longevidade	**10-20 anos**
Ativo	**De dia**
Estilo de vida	**Terrestre e escavador**
Habitat	**Semideserto e savana**
Dificuldade	**Especialista**

CUIDADOS

Terrário mín.	**300 × 200 × 150 cm** C × L × A para 1 adulto
Temperatura, dia	**24-26 °C**
Ponto quente	**35-45 °C**
Temperatura, noite	**19-21 °C**
Umidade	**30-60%**
Água	**Recipiente de água**
UVB	**Recomendado** Zona de Ferguson 3
Alimentação	**Insetos, vertebrados e minhocas/caracóis**
Mantido	**Sozinho ou em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Tamanho da postura	**16-65 ovos**
Posturas por ano	**1-2**
Incubação	**150-180 dias** a 28-31 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
Anexo da UE	**Anexo B**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO

An adult *Varanus albigularis* needs at least 300 × 200 × 150 cm (length × width × height) and preferably a whole room.

TEMPERATURA

As a diurnal basker, *Varanus albigularis* requires a large temperature gradient. The air temperature is 24-26 °C by day and 19-21 °C by night.

UMIDADE E ÁGUA

The species comes from dry to semi-humid savanna country, and the right humidity in the setup is not documented with figures.

ILUMINAÇÃO E UVB

Varanus albigularis is a diurnal sun-basker and needs strong light.

ALIMENTAÇÃO

In the wild *Varanus albigularis* is a true generalist. It eats snails, grasshoppers, millipedes, beetles and other invertebrates and occasionally snakes.

COMPORTAMENTO E MANUSEIO

Varanus albigularis is not a pet in the usual sense. Wild animals are shy and flee or bite if they are caught, and even tame animals have strong jaws, sharp claws and a tail that can lash.

ÁREA DE OCORRÊNCIA NATURAL

Angola, Botsuana, Etiópia, Quênia, Lesoto, Malaui, Moçambique, Namíbia, Somália, Essuatíni, Tanzânia, Uganda, África do Sul, Zâmbia, Zimbábue e Congo - Kinshasa.