herpy

VARANOS

Spotted Tree Monitor

Varanus timorensis

Varanus timorensis is a small, diurnal tree monitor from Timor and the neighbouring islands of the Lesser Sunda Islands, dark with yellow spots and a very long tail. It grows to no more than 61 cm, but needs a tall, warm enclosure, daily feeding with live insects and a keeper with experience of small monitors.

Família: Varanidae Nome anterior: *Varanus auffenbergi*

DADOS PRINCIPAIS

O ANIMAL

Tamanho adulto	**50-61 cm** comprimento total com a cauda
Peso	**100-350 g**
Longevidade	**10-15 anos**
Ativo	**De dia**
Estilo de vida	**Arborícola**
Habitat	**Floresta seca e floresta tropical**
Dificuldade	**Experiente**

CUIDADOS

Terrário mín.	**150 × 75 × 150 cm** C × L × A para 1 adulto
Temperatura, dia	**27-32 °C**
Ponto quente	**40-50 °C**
Temperatura, noite	**22-25 °C**
Umidade	**50-70%**
Água	**Recipiente de água**
UVB	**Recomendado** Zona de Ferguson 3
Alimentação	**Insetos e vertebrados**
Mantido	**Sozinho ou em casal**

REPRODUÇÃO

Reprodução	**Ovíparo**
Tamanho da postura	**5-11 ovos**
Posturas por ano	**1-2**
Incubação	**90-135 dias** a 29-32 °C

PROTEÇÃO E DOCUMENTAÇÃO

CITES	**Apêndice II**
Anexo da UE	**Anexo B**
IUCN	**Menos preocupante (LC)**

Resumo dos cuidados

O TERRÁRIO	An adult needs a tall enclosure of at least 150 × 75 × 150 cm (length × width × height). Larger is better, because the species is active and needs space for both a warm and a cool zone.
TEMPERATURA	*Varanus timorensis* needs 27-32 °C in the air by day and 22-25 °C at night.
UMIDADE E ÁGUA	Humidity in the enclosure should be 50-70 %, and the enclosure should be well ventilated.
ILUMINAÇÃO E UVB	As a diurnal sun-lover, *Varanus timorensis* needs strong light and UVB all day.
ALIMENTAÇÃO	In the wild *Varanus timorensis* feeds mostly on arthropods, but also takes lizards and small snakes.
COMPORTAMENTO E MANUSEIO	*Varanus timorensis* is a small, quick animal that flees into hiding when startled. Bites from an animal of up to 61 cm are rarely life-threatening, but the teeth are sharp, and the claws can tear the skin.
ÁREA DE OCORRÊNCIA NATURAL	Indonésia e Timor-Leste.